

August 15, 1960

THE WILCOXON TWO SAMPLE STATISTIC

James E. Jacobson<sup>1</sup>/

Special Report No. 2

Department of Statistics  
University of Minnesota  
Minneapolis, Minnesota

Reproduction in whole or in part is permitted for any  
purpose of the United States Government

<sup>1</sup>/ Work done in part under contract Nonr2582(00), Task  
NR 042-200 of the Office of Naval Research

# THE WILCOXON TWO SAMPLE STATISTIC

James E. Jacobson

## ERRATA SHEET

<u>Page</u>	<u>Line</u>	<u>Comments</u>
1	6	Lower case "o" to upper case "O".
1	16	Period to comma and upper case "O" to lower case "o".
1	18	"all real" to "some".
1	25	Omit "the distribution of".
1	27	"parenthesis" to "parentheses".
4	26	$\delta = \alpha$ to $\delta = \gamma$
4	27	$\delta = \gamma$ to $\alpha = \gamma$
5		Substitute Chart I (revised)
6	9	$\alpha$ to .025
6	13	$\alpha$ to .05
7	19	$\begin{array}{ccccc} 9 & 0 & +1 & \text{to} & 9 & 0 & +1 \\ & \frac{1}{2} & +1/0 & & & \frac{1}{2} & +1/0^c \end{array}$
7	42	Add "(or additional continuity correction)"
8	2	Lower case "t" to upper case "T".
17	1	$\delta = .005$ to $\delta = \gamma$
33	40	Annals of Eugenics to <u>Annals of Eugenics</u> .
34	9	"l;egalite" to "l'egalite".
36	35	7(1952) to 8(1952).
37	4	"Stanford" to "Stanford".
37	5	"Stanford" to "Stanford".

Table II gives values of  $\delta = \gamma$ , not  $\delta = \alpha$ .

Table III gives values of  $\alpha = \gamma$ , not  $\delta = \gamma$ .

# The Wilcoxon Two Sample Statistic

## 1. Summary.

This paper presents tables for the Wilcoxon two sample statistic and considers the normal approximation to the Wilcoxon two sample statistic. The primary purposes of this paper are a compilation of existing tables of the exact distribution of the Wilcoxon statistic and a comparison of the exact distribution ( in the tails ) to the normal approximation. Of secondary importance are some additional exact percentage points computed for the Wilcoxon statistic. The remainder of this paper is divided into the following sections: (2) Wilcoxon two sample statistic, (3) Existing tables of the exact distribution of the statistic, (4) Description of tables presented here, (5) Normal approximation, (6) Continuity correction, (7) Computational method, (8) Tables, and (9) Bibliography.

## 2. Wilcoxon two sample statistic.

Given two independent random samples  $X_1, \dots, X_m$  and  $Y_1, \dots, Y_n$  from two populations with unknown cumulative distribution functions,  $F$  and  $G$ , respectively, One can test the hypothesis  $H_0: F(t) = G(t)$  against the one-sided alternative hypothesis  $H_A: F(t) \geq G(t)$  and  $F(t) \neq G(t)$  for ~~all~~<sup>some</sup> real  $t$ . Wilcoxon (55)<sup>1</sup> proposed the statistic  $\sum_{i=1}^m r_i$  where  $r_1, \dots, r_m$  are the  $X$  ranks in the combined sample.

Equivalent to the Wilcoxon two sample statistic is  $U$ , the Mann-Whitney statistic, the number of pairs  $(X_i, Y_j)$  for which  $Y_j < X_i$ . For the one-sided test the rejection region is  $U \leq t$  where  $t = t_{\alpha, m, n}$  is the greatest integer for which  $P(U \leq t | H_0) \leq \alpha$ , the desired level of significance. Under  $H_0$  Mann and Whitney (27) proved that ~~the distribution of~~  $U$  is asymptotically normally distributed when  $m$  and  $n$  approach infinity. Under  $H_0$  the mean

1. The number in the paratheses refers to the listed references in the bibliography.

and the variance of  $U$  are  $mn/2$  and  $mn(m+n+1)/12$ , respectively.

### 3. Existing tables of the exact distribution of the statistic.

Chart I shows the extent of existing tables and the extent of the tables presented here.

The present tables are compared with the previous as follows:

- a) Mann-Whitney's tables (27) for  $\alpha = .005, .01, .025, .05$  over the range  $\text{Max.}(m,n) \leq 8$ .
- b) van der Vaart's tables (46) for  $\alpha = .005, .01, .025, .05$  over the range  $\text{Max.}(m,n) \leq 10$ .
- c) Auble's tables (1) for  $\alpha = .005, .01, .025, .05$  over the range  $\text{Min.}(m,n) \leq 12$  and  $\text{Max.}(m,n) \leq 20$ .
- d) Natrella's tables<sup>2</sup> (29) for  $\alpha = .005, .01, .025$  over the range  $\text{Min.}(m,n) \leq 12$  and  $m+n \leq 30$ .

Let  $t$  be the largest integer such that  $P(U \leq t | H_0) \leq \alpha$ . Then the differences with the previous tables are as follows:

for  $\alpha = .005$

$m = 7, n = 5$

van der Vaart (46)  $t = 2$

given value  $t = 1$

$m = 16, n = 13$

Natrella (29)  $t = 46$

given value  $t = 45$

$m = 21, n = 5$

Natrella (29)  $t = 14$

given value  $t = 13$

2. Since Natrella's tables encompass White's (52) for the part compared here of the two only Natrella's is checked.

for  $\alpha = .01$

$m = 8, n = 7$

Mann-Whitney (27)

$t = 8^3$

given value

$t = 7$

$m = 8, n = 8$

Mann-Whitney (27)

$t = 10^4$

given value

$t = 9$

$m = 18, n = 9$

Natrella (29)

$t = 35$

given value

$t = 36$

for  $\alpha = .025$

$m = 11, n = 2$

Natrella (29)

$t = 1$

given value

$t = 0$

$m = 13, n = 13$

Natrella (29)

$t = 46$

given value

$t = 45$

$m = 15, n = 15$

Natrella (29)

$t = 65$

given value

$t = 64$

$m = 16, n = 11$

Natrella (29)

$t = 48$

given value

$t = 47$

$m = 17, n = 14$

Auble (1)

$t = 67$

given value

$t = 69$

3. Mann-Whitney gives  $P(U \leq 8 | H_0) = .010$  to three places, but to six places is .010256.

4. Mann-Whitney gives  $P(U \leq 10 | H_0) = .010$  to three places, but to six places is .010335.

$m = 20, n = 4$

Auble (1)  $t = 13$

given value  $t = 14$

$m = 21, n = 9$

Natrella (29)  $t = 50$

given value  $t = 51$

for  $\alpha = .05$

$m = 9, n = 3$

Auble (1)  $t = 3$

given value  $t = 4$

$m = 19, n = 7$

Auble (1)  $t = 27^5$

given value  $t = 37$


#### 4. Description of tables presented here.

Working under the null hypothesis,  $H_0$ , and considering only the lower tail of the distribution, the significant values of the Wilcoxon two sample test and the normal approximation to the Wilcoxon two sample probability distribution have been computed and compared for the levels of significance of 0.5, 1, 2.5, and 5 per cent, and those  $m$  and  $n$  for which  $\text{Min.}(m, n) \leq 12$  and  $m + n \leq 30$ . Table I shows the largest integer,  $t$ , for which  $\gamma = P(U \leq t | H_0) \leq \alpha$ , the desired level of significance, and the corresponding normal approximation. (The  $t$ 's corresponding to the exact values outside the range of my computation -  $\text{Min.}(m, n) \leq 12$  and  $m + n \leq 30$  - are taken from Auble (1).) Table II shows  $\delta - \alpha$  where  $\delta = P(U \leq t + 1 | H_0) > \alpha$ . Table III shows  $\alpha - \gamma$ .

#### 5. Typographical error.

Chart I

Existing tables of the Wilcoxon two sample statistic and levels of significance for the one-sided test.



## 5. Normal approximation.

The notation and restrictions of section 4 will be used here. At the .005 level of significance the normal approximation is  $t$  or  $t - 1$  for the minimum of sample sizes greater than 3. At the .01 level of significance the normal approximation is  $t$  ( most often ) or  $t + 1$  for the minimum of the two sample sizes greater than 3. At the .025 level of significance the normal approximation is  $t$  and  $t + 1$  ( most often ). Hence for the sample sizes considered using the normal approximation minus one will yield a level of significance less than or equal to  $\alpha$ <sup>.025</sup>. At the .05 level of significance the normal approximation is  $t + 1$  or  $t + 2$  ( most often ). Hence for the sample sizes considered using the normal approximation minus two will yield a level of significance less than or equal to  $\alpha$ <sup>.05</sup>.

## 6. Continuity correction.

Among the possible combinations of continuity correction and rounding rule obtained by using a continuity correction,  $x$ , which can be represented as a fraction with denominator 2 and using a rounding rule of round up, round down, or round according to the usual rule there are two distinct sets with respect to the differences between the normal approximation and the exact values for the Wilcoxon two sample test. A continuity correction of  $x$  and round down,  $x + \frac{1}{2}$  and round according to the usual rule, or  $x + 1$  and round up give the same result. Substituting  $x - 1$  or  $x + 1$  for  $x$  in the previous statement will yield one more or one less, respectively, than the previous result. Changing  $x$  by one half plus an integer yields the second set. Table A shows the number of differences between the normal approximation and the exact value of the Wilcoxon two sample test for continuity corrections



Table A

The Number of Differences Between the Exact Values of the Wilcoxon Test and the Normal Approximation for  $1 \leq n \leq m \leq 20$ .

levels of significance									
	continuity m correction	$\alpha^{.05}$		$\alpha^{.025}$		$\alpha^{.01}$		$\alpha^{.005}$	
		C.F.	D/T <sup>b</sup>	C.F.	D/T	C.F.	D/T	C.F.	D/T
3	0 $\frac{1}{2}$	+1 0	0/1 0/1	- -	- -	- -	- -	- -	- -
4	0 $\frac{1}{2}$	+1 +1	0/2 0/2	+1 +1	0/1 0/1	- -	- -	- -	- -
5	0 $\frac{1}{2}$	+1 0	0/4 1/4	+1 0	0/3 1/3	+1/0 <sup>c</sup> 0	1/2 0/2	0 0	0/1 0/1
6	0 $\frac{1}{2}$	+1 0	0/5 2/5	+1 0	1/4 1/4	0 0	0/3 0/3	0 +1	0/3 0/3
7	0 $\frac{1}{2}$	+1 +1	0/6 1/6	+1 0	1/5 2/5	0 0	2/5 0/5	0 0	1/4 1/4
8	0 $\frac{1}{2}$	+1 0	0/7 3/7	0 0	3/7 1/7	+1/0 0	3/6 1/6	0 0	0/5 2/5
9	0 $\frac{1}{2}$	+1 +1/0 <sup>c</sup>	0/8 4/8	+1 0	1/8 2/8	0 0	2/7 2/7	0 0	0/7 1/7
10	0 $\frac{1}{2}$	+1 +1	0/9 3/9	+1 0	1/9 2/9	+1/0 0	4/8 0/8	0 -1	0/8 3/8
11	0 $\frac{1}{2}$	+1 +1/0	1/10 5/10	+1 0	1/10 4/10	+1 0	4/9 2/9	0 -1	1/9 4/9
12	0 $\frac{1}{2}$	+1 +1	3/11 3/11	+1 +1	4/11 4/11	0 0	3/10 3/10	0 -1	2/10 2/10
13	0 $\frac{1}{2}$	+1 +1	0/12 3/12	+1 0	2/12 2/12	0 0	2/12 3/12	0 -1	4/11 3/11
14	0 $\frac{1}{2}$	+1 +1	2/13 4/13	+1 0	2/13 3/13	0 0	6/13 3/13	0 -1	1/12 4/12
15	0 $\frac{1}{2}$	+1 0	3/14 6/14	+1 0	2/14 3/14	0 0	3/14 5/14	0 -1	3/13 4/13
16	0 $\frac{1}{2}$	+1 +1	0/15 6/15	+1 0	6/15 3/15	0 0	1/15 7/15	0 -1	3/14 5/14
17	0 $\frac{1}{2}$	+1 0	4/16 7/16	+1 0	3/16 3/16	0 0/-1	2/16 8/16	0 -1	5/15 1/15
18	0 $\frac{1}{2}$	+1 +1	2/17 7/17	+1 0	4/17 4/17	0 0	4/17 7/17	0 -1	6/16 2/16
19	0 $\frac{1}{2}$	+1 +1	1/19 6/19	+1 0	4/18 4/18	0 0	2/18 5/18	-1 -1	8/18 8/18
20	0 $\frac{1}{2}$	+1 0	1/20 9/20	+1 0	3/19 3/19	0 0	5/19 8/19	-1 -1	9/19 5/19

a. correction factor, b. difference/total, c. +1 or 0.  
(or additional continuity correction)

of 0 and  $\frac{1}{2}$  and round according to the usual rule and for  $1 \leq n \leq m \leq 20$ . Checking Table A a continuity correction of 1 and round according the usual rule is best for levels of significance .05 and .025 and a continuity correction of 0 and round according to the usual rule is best for levels of significance .01 and .005.

#### 7. Computational method.

The exact probabilities for the Wilcoxon two sample test were computed using the method - a difference equation in three variables and two auxiliary tables - shown by Fix and Hodges (11). Computations were continued as far as practicable with a desk computer which is dependent on the extent of the auxiliary tables given by Fix and Hodges - i.e.,  $\text{Min.}(m,n) \leq 12$ . For a variety of methods that may be used to compute the exact probabilities see van der Vaart (48). In table I the normal approximation was computed using the property that  $\frac{U - \frac{1}{2}mn}{\sqrt{mn(m+n+1)/12}}$  approaches  $N(0,1)$  as  $m$  and  $n$  approach infinity. No continuity correction was used and rounding was according to the usual rule.

Table I ( $\alpha = .005$ )

9

Exact values for the .005 level of significance and the corresponding normal approximation for  $\text{Max.}(m,n) \leq 20$  or  $m + n \leq 30$ .

m-n	1		2		3		4		5		6		7		8	
	E <sup>a</sup>	N <sup>b</sup>	E	N	E	N	E	N	E	N	E	N	E	N	E	N
0	-	-	-	-	-	-	-	-	0	0	2	2	4	4	7	7
1	-	-	-	-	-	-	-	-	1	1	3	3	6	6	9	9
2	-	-	-	-	-	-	0	0	1	2	4	4	7	7	11	11
3	-	-	-	-	-	-	0	0	2	2	5	5	9	9	13	13
4	-	-	-	-	-	-	1	1	3	3	6	6	10	10	15	15
5	-	-	-	-	-	-	1	1	4	4	7	7	12	12	17	16
6	-	-	-	-	0	0	2	2	5	5	9	8	13	13	18	18
7	-	-	-	-	0	0	2	2	6	6	10	10	15	14	20	20
8	-	-	-	-	0	0	3	3	7	6	11	11	16	16	22	22
9	-	-	-	-	1	0	3	3	7	7	12	12	18	17	24	24
10	-	-	-	-	1	0	4	4	8	8	13	13	19	19	26	26
11	-	-	-	-	1	1	5	4	9	9	15	14	21	20	28	27
12	-	-	-	-	2	1	5	5	10	10	16	15	22	22	30	29
13	-	-	-	-	2	1	6	5	11	10	17	17	24	23	32	31
14	-	-	-	-	2	1	6	6	12	11	18	18	25	25	34	33
15	-	-	-	-	2	1	7	6	13	12	19	19	27	26		
16	-	-	-	-	3	2	8	7	13	13	21	20	29	28		
17	-	-	0	-3	3	2	8	7	14	14	22	21				
18	-	-	0	-3	3	2	9	8	15	15	23	22				
19	-	-	0	-3	4	2	9	8	16	15						
20	-	-	0	-3	4	2	10	9	17	16						
21	-	-	0	-3	4	3	10	9								
22	-	-	0	-3	5	3	11	10								
23	-	-	0	-3	5	3										
24	-	-	0	-3	5	3										
25	-	-	1	-3												
26	-	-	1	-3												

a. exact value

b. normal approximation

Table I ( $\alpha = .005$ )

10

m-n	9		10		11		12		13		14		15		16	
	E	N	E	N	E	N	E	N	E	N	E	N	E	N	E	N
0	11	11	16	16	21	20	27	27	34	34	42	42	51	50	60	60
1	13	13	18	18	24	24	31	31	38	38	46	46	55	55	65	64
2	16	16	21	21	27	27	34	34	42	42	50	50	60	59	70	69
3	18	18	24	23	30	30	37	37	45	45	54	54	64	64	74	74
4	20	20	26	26	33	33	41	41	49	49	58	58	68	68	79	79
5	22	22	29	29	36	36	44	44	53	53	63	62	73	73		
6	24	24	31	31	39	39	47	47	56	56	67	66				
7	27	26	34	34	42	42	51	50	60	60						
8	29	29	37	36	45	45	54	54								
9	31	31	39	39	48	48										
10	33	33	42	41												
11	36	35														
12	38	38														

m-n	17		18		19		20	
	E	N	E	N	E	N	E	N
0	70	70	81	81	93	92	105	105
1	75	75	87	86	99	98		
2	81	80	92	92				
3	86	85						

limit value for normal approximation as $m \rightarrow \infty$	n			
	1	2	3	4
	-0.244	-0.052	0.212	0.513

11

III-14

[illegible]

Table I ( $\alpha = .01$ )

12

m-n	9		10		11		12		13		14		15		16	
	E	N	E	N	E	N	E	N	E	N	E	N	E	N	E	N
0	14	14	19	19	25	25	31	32	39	39	47	47	56	56	66	66
1	16	17	22	22	28	28	35	35	43	43	51	52	61	61	71	71
2	18	19	24	25	31	31	38	39	47	47	56	56	66	66	76	77
3	21	21	27	27	34	35	42	42	51	51	60	60	70	71	82	82
4	23	24	30	30	37	38	46	46	55	55	65	65	75	75	87	87
5	26	26	33	33	41	41	49	49	59	59	69	69	80	80		
6	28	28	36	36	44	44	53	53	63	63	73	74				
7	31	31	38	39	47	47	56	57	67	67						
8	33	33	41	41	50	50	60	60								
9	36	36	44	44	53	54										
10	38	38	47	47												
11	40	41														
12	43	43														

m-n	17		18		19		20	
	E	N	E	N	E	N	E	N
0	77	77	88	88	101	101	114	114
1	82	83	94	94	107	107		
2	88	88	100	100				
3	93	94						

limit value for normal approximation as $m \rightarrow \infty$	1		n		3		4	
	1	2	2	3	3	4	4	5
	-0.171	0.050	0.337	0.657				

III-11

[illegible]

Table I ( $\alpha = .025$ )

14

m-n	9		10		11		12		13		14		15		16	
	E	N	E	N	E	N	E	N	E	N	E	N	E	N	E	N
0	17	18	23	24	30	31	37	38	45	46	55	55	64	65	75	76
1	20	21	26	27	33	34	41	42	50	51	59	60	70	70	81	82
2	23	24	29	30	37	38	45	46	54	55	64	65	75	76	86	87
3	26	26	33	33	40	41	49	50	59	59	69	70	80	81	92	93
4	28	29	36	37	44	45	53	54	63	64	74	74	85	86	98	98
5	31	32	39	40	47	48	57	58	67	68	78	79	90	91		
6	34	35	42	43	51	52	61	62	72	72	83	84				
7	37	37	45	46	55	55	65	66	76	77						
8	39	40	48	49	58	59	69	70								
9	42	43	52	52	62	63										
10	45	46	55	55												
11	48	48														
12	51	51														

m-n	17		18		19		20	
	E	N	E	N	E	N	E	N
0	87	88	99	100	113	113	127	128
1	93	94	106	106	119	120		
2	99	100	112	113				
3	105	106						

limit value for normal approximation as $m \rightarrow \infty$	1		2		3		4	
	-0.066		0.200		0.520		0.868	



m-n

[illegible]

Table I ( $\alpha = .05$ )

16

m-n	9		10		11		12		13		14		15		16	
	E	N	E	N	E	N	E	N	E	N	E	N	E	N	E	N
0	21	22	27	28	34	35	42	44	51	52	61	62	72	73	83	84
1	24	25	31	32	38	39	47	48	56	57	66	67	77	78	89	90
2	27	28	34	35	42	43	51	52	61	62	71	72	83	84	95	96
3	30	31	37	38	46	47	55	56	65	66	77	78	88	90	101	102
4	33	34	41	42	50	51	60	61	70	71	82	83	94	95	107	108
5	36	37	44	45	54	55	64	65	75	76	87	88	100	101		
6	39	40	48	49	57	59	68	69	80	81	92	93				
7	42	43	51	52	61	62	72	73	84	85						
8	45	46	55	56	65	66	77	78								
9	48	49	58	59	69	70										
10	51	52	62	63												
11	54	55														
12	57	58														

m-n	17		18		19		20	
	E	N	E	N	E	N	E	N
0	96	97	109	110	123	124	138	139
1	102	103	116	117	130	131		
2	109	110	123	124				
3	115	116						

m-n	1		2		3		4	
	E	N	E	N	E	N	E	N
27	0	0						
28	0	0						
limit value for normal approximation as $m \rightarrow \infty$	0.025		0.028		0.678		1.050	

Table II ( $\alpha = .005$ )  
 $\delta = .005$  where  $t + 1$  is the smallest integer such that  
 $\delta = P(U \leq t + 1 | H_0) > .005$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

m-n	1	2	3	4	5	6	7	8
0	-	-	-	-	.003968	.003247	.002040	.001709
1	-	-	-	-	.004329	.002914	.002331	.001604
2	-	-	-	.004762	.002525	.002331	.001836	.001554
3	-	-	-	.003030	.002331	.002198	.001954	.001482
4	-	-	-	.004040	.002498	.001748	.001540	.001421
5	-	-	-	.002797	.002331	.001616	.001608	.001356
6	-	-	.004545	.002997	.002289	.001885	.001342	.001063
7	-	-	.003497	.002198	.002101	.001622	.001393	.001032
8	-	-	.002747	.002747	.002101	.001496	.001167	.000998
9	-	-	.004396	.002101	.001548	.001308	.001207	.000966
10	-	-	.003571	.001961	.001483	.001206	.001040	.000935
11	-	-	.002941	.002322	.001474	.001347	.001071	.000905
12	-	-	.003676	.001858	.001405	.001211	.000929	.000877
13	-	-	.003096	.001838	.001397	.001124	.000955	.000850
14	-	-	.002632	.001504	.001341	.001021	.000842	.000824
15	-	-	.002256	.001694	.001318	.000953	.000864	
16	-	-	.002597	.001694	.001277	.001038	.000909	
17	-	.004762	.002259	.001423	.001041	.000956		
18	-	.004329	.001976	.001538	.001028	.000896		
19	-	.003953	.002174	.001311	.001002			
20	-	.003623	.001923	.001319	.000989			
21	-	.003333	.001709	.001137				
22	-	.003077	.002137	.001241				
23	-	.002849	.001916					
24	-	.002646	.001724					
25	-	.004926						
26	-	.004598						

Table II ( $\alpha = .005$ )

m-n	n			
	9	10	11	12
0	.001419	.001283	.001026	.000876
1	.001256	.001072	.000974	.000919
2	.001387	.001098	.000929	.000826
3	.001235	.001104	.000882	.000749
4	.001112	.000942	.000841	.000776
5	.001003	.000951	.000801	.000707
6	.000914	.000823	.000764	.000643
7	.000982	.000827	.000730	.000665
8	.000897	.000830	.000652	.000612
9	.000825	.000731	.000669	
10	.000759	.000732		
11	.000802			
12	.000742			

Table II ( $\alpha = .01$ )

$\delta = .01$  where  $t + 1$  is the smallest integer such that  
 $\delta = P(U \leq t + 1 | H_0) > .01$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

$m-n$	1	2	3	4	5	6	7	8
0	-	-	-	-	.007937	.005411	.004371	.002953
1	-	-	-	.007937	.006494	.004079	.003263	.002715
2	-	-	-	.009524	.006313	.004662	.003234	.002491
3	-	-	-	.006061	.005439	.003996	.003239	.002302
4	-	-	.008333	.006061	.004995	.003247	.002514	.002135
5	-	-	.006061	.006993	.004329	.002828	.002520	.002359
6	-	-	.009091	.004995	.004121	.003124	.002503	.002183
7	-	-	.006993	.004396	.003717	.002617	.002073	.002031
8	-	-	.005595	.004945	.003501	.002322	.002064	.001897
9	-	-	.006593	.003782	.003182	.002488	.002052	.001778
10	-	-	.005357	.003595	.003031	.002171	.002031	.001671
11	-	.009524	.004412	.003870	.002801	.002308	.001727	.001577
12	-	.008333	.004902	.003096	.002658	.002080	.001719	.001491
13	-	.007353	.004128	.003008	.002496	.001858	.001706	.001413
14	-	.006536	.004386	.003144	.002376	.001950	.001692	.001343
15	-	.005848	.003759	.002597	.002240	.001784	.001480	
16	-	.005263	.003247	.002541	.002144	.001614	.001471	
17	-	.009524	.003953	.002688	.002031	.001593		
18	-	.008658	.003458	.002274	.001954	.001556		
19	-	.007905	.003478	.002222	.001861			
20	-	.007246	.003077	.002295	.001789			
21	-	.006667	.002735	.001979				
22	-	.006154	.003053	.001970				
23	-	.005698	.002737					
24	-	.005291	.002463					
25	-	.004926						
26	-	.004598						

Table II ( $\alpha = .01$ )

m-n	n			
	9	10	11	12
0	.002838	.002344	.002038	.001609
1	.002403	.002245	.001853	.001610
2	.002072	.001840	.001699	.001413
3	.002164	.001794	.001561	.001411
4	.001890	.001735	.001444	.001404
5	.001942	.001684	.001512	.001253
6	.001717	.001628	.001401	.001246
7	.001761	.001396	.001304	.001122
8	.001572	.001362	.001218	.001117
9	.001605	.001325	.001141	
10	.001445	.001291		
11	.001311			
12	.001336			

$\delta = .025$  where  $t + 1$  is the smallest integer such that  
 $\delta = P(U \leq t + 1 | H_0) > .025$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

m-n	1	2	3	4	5	6	7	8
0	-	-	-	.014286	.011905	.011905	.008159	.007537
1	-	-	-	.015873	.010823	.008159	.006993	.006417
2	-	-	.017857	.014286	.012626	.008325	.006556	.005533
3	-	-	.023810	.015152	.010101	.008392	.006119	.004869
4	-	-	.016667	.012121	.008991	.006993	.005719	.005017
5	-	-	.018182	.008392	.007659	.006949	.005358	.004467
6	-	.022222	.013636	.008991	.006868	.005710	.005057	.004006
7	-	.018182	.013986	.008059	.007595	.005750	.004764	.004138
8	-	.015152	.010989	.008242	.006653	.004954	.004515	.003745
9	-	.012821	.010989	.007563	.006020	.004976	.003467	.003825
10	-	.021978	.008921	.007516	.005418	.004986	.004074	.003490
11	-	.019048	.010294	.006966	.004963	.004309	.003882	.003203
12	-	.016667	.008578	.007018	.005316	.004331	.003709	.003273
13	-	.014706	.008256	.005681	.004844	.003840	.003548	.003021
14	-	.013072	.007018	.005332	.004494	.003861	.003402	.002800
15	-	.011696	.007519	.005308	.004141	.003871	.003266	
16	-	.010526	.006494	.005082	.004393	.003451	.003140	
17	-	.009524	.006776	.005059	.004100	.003465		
18	-	.008658	.005928	.004816	.003816	.003136		
19	-	.011858	.006087	.004786	.003579			
20	-	.010870	.005385	.004103	.003761			
21	-	.010000	.005470	.003958				
22	-	.009231	.004884	.003941				
23	-	.008547	.005200					
24	-	.007937	.004680					
25	-	.007389						
26	-	.006897						

m-n	n			
	9	10	11	12
0	.005162	.004584	.004180	.003481
1	.005001	.004190	.003670	.003306
2	.004870	.003877	.003651	.003145
3	.004705	.004064	.003244	.002996
4	.003989	.003754	.003227	.002859
5	.003893	.003493	.002902	.002732
6	.003790	.003257	.002887	.002616
7	.003691	.003053	.002864	.002508
8	.003228	.002867	.002606	.002408
9	.003161	.002961	.002588	
10	.003087	.002791		
11	.003017			
12	.002696			



Table II ( $\alpha = .05$ )

$\delta = .05$  where  $t + 1$  is the smallest integer such that  
 $\delta = P(U \leq t + 1 | H_0) > .05$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

m-n	1	2	3	4	5	6	7	8
0	-	-	.050000	.028571	.027778	.019481	.015443	.010956
1	-	-	.028571	.023810	.021645	.015486	.013364	.010613
2	-	-	.035714	.023810	.016414	.013320	.011713	.008981
3	-	.047619	.035714	.018182	.017094	.012787	.010490	.008904
4	-	.035714	.025000	.018182	.014486	.012238	.009427	.008693
5	-	.027778	.024242	.015385	.014985	.011635	.008573	.007573
6	-	.044444	.022727	.014985	.012592	.009696	.008914	.007480
7	-	.036364	.017483	.013158	.010989	.009325	.008153	.006612
8	-	.030303	.019231	.012637	.011321	.009004	.007511	.006518
9	-	.025641	.015385	.011345	.009890	.008698	.006955	.006434
10	-	.021978	.014286	.011111	.010126	.008391	.007183	.005788
11	-	.019048	.014706	.010062	.009091	.007930	.006686	.005712
12	-	.025000	.012255	.011146	.008126	.007014	.006242	.005641
13	-	.022059	.012384	.010693	.008291	.006832	.005855	.005138
14	-	.019608	.012281	.009843	.007552	.006654	.005507	.005077
15	-	.017544	.010526	.009486	.007698	.006480	.005644	
16	-	.015789	.010390	.008846	.007023	.006312	.005327	
17	-	.014286	.010728	.008538	.007135	.005652		
18	.050000	.012987	.009387	.008027	.006583	.005531		
19	.047619	.015810	.009130	.007749	.006071			
20	.045455	.014493	.009230	.007326	.006175			
21	.043478	.013333	.008205	.007115				
22	.041667	.012308	.008242	.006751				
23	.040000	.011396	.008210					
24	.038462	.010582	.007389					
25	.037037	.012315						
26	.035714	.011494						

Table II ( $\alpha = .05$ )

m-n	n			
	9	10	11	12
0	.009790	.007956	.006735	.005898
1	.009017	.007924	.006433	.005953
2	.008383	.007056	.006153	.005494
3	.007801	.006359	.005891	.005095
4	.007312	.006351	.005648	.005119
5	.006864	.005768	.005420	.004773
6	.006475	.005752	.004825	.004467
7	.006119	.005277	.004660	.004195
8	.005804	.005259	.004505	.004212
9	.005523	.004855	.004358	
10	.005256	.004838		
11	.005017			
12	.004800			

m-n	n
	1
27	.034483
28	.033333

.005

Table III ( $\alpha = .005$ )

$\delta = r$  where  $t$  is the largest integer such that  $\gamma = P(U \leq t | H_0) \leq .005$   
 and  $\delta = P(U \leq t + 1 | H_0)$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

$m-n$	1	2	3	4	5	6	7	8
0	-	-	-	-	.001032	.000671	.001503	.001503
1	-	-	-	-	.000671	.000921	.000338	.001051
2	-	-	-	.000238	.002475	.001004	.001066	.000726
3	-	-	-	.001970	.001892	.001204	.000167	.000449
4	-	-	-	.000960	.001503	.001254	.000852	.000221
5	-	-	-	.002203	.001004	.001445	.000118	.000032
6	-	-	.000455	.001004	.000660	.000152	.000769	.000938
7	-	-	.001503	.002070	.000314	.000393	.000158	.000772
8	-	-	.002253	.001154	.000098	.000640	.000737	.000626
9	-	-	.000604	.002059	.001388	.000817	.000211	.000497
10	-	-	.001429	.001078	.001130	.001006	.000741	.000384
11	-	-	.002059	.000356	.000921	.000067	.000278	.000282
12	-	-	.000098	.001285	.000709	.000290	.000756	.000191
13	-	-	.000872	.000489	.000542	.000500	.000342	.000110
14	-	-	.001491	.001309	.000365	.000674	.000781	.000036
15	-	-	.001992	.000709	.000219	.000841	.000407	
16	-	-	.000455	.000012	.000074	.000105	.000041	
17	-	.000238	.001047	.000810	.000987	.000295		
18	-	.000671	.001542	.000251	.000849	.000471		
19	-	.001047	.000217	.000954	.000714			
20	-	.001377	.000769	.000409	.000593			
21	-	.001667	.001239	.001042				
22	-	.001923	.000116	.000584				
23	-	.002151	.000621					
24	-	.002354	.001059					
25	-	.000074						
26	-	.000402						

Table III ( $\alpha = .005$ )

m-n	n			
	9	10	11	12
0	.001113	.000535	.000825	.000853
1	.001190	.001025	.000687	.000222
2	.000154	.000522	.000576	.000443
3	.000356	.000056	.000489	.000645
4	.000545	.000574	.000423	.000146
5	.000718	.000192	.000372	.000371
6	.000872	.000657	.000334	.000573
7	.000179	.000330	.000306	.000163
8	.000376	.000022	.000285	.000375
9	.000547	.000463	.000271	
10	.000714	.000192		
11	.000156			
12	.000338			

Table III ( $\alpha = .01$ )								
$\delta = \gamma$ where $t$ is the largest integer such that $\gamma = P(U \leq t   H_0) \leq .01$ and $\delta = P(U \leq t + 1   H_0)$ for $m + n \leq 30$ and $\text{Min.}(m, n) \leq 12$ .								
$m-n$	1	2	3	4	5	6	7	8
0	-	-	-	-	.002063	.002424	.001259	.002618
1	-	-	-	.002063	.001342	.003007	.003007	.002390
2	-	-	-	.000471	.001162	.000010	.001783	.002230
3	-	-	-	.003939	.000676	.001209	.000744	.002128
4	-	-	.001667	.001919	.000509	.002008	.002301	.003070
5	-	-	.003939	.000210	.000343	.002728	.001466	.000054
6	-	-	.000909	.003007	.000385	.000896	.000699	.000174
7	-	-	.003007	.001208	.000304	.001633	.002071	.000294
8	-	-	.004405	.000110	.000313	.002312	.001422	.000412
9	-	-	.001209	.002437	.000282	.000823	.000818	.000525
10	-	-	.002857	.001176	.000325	.001503	.000257	.000634
11	-	.000476	.004118	.000196	.000319	.000143	.001460	.000738
12	-	.001667	.001422	.002157	.000355	.000862	.000964	.000837
13	-	.002647	.002776	.001245	.000371	.001462	.000494	.000931
14	-	.003464	.000351	.000294	.000401	.000305	.000050	.001021
15	-	.004152	.001729	.001982	.000420	.000926	.001111	
16	-	.004737	.002857	.001154	.000453	.001458	.000705	
17	-	.000476	.000966	.000435	.000474	.000453		
18	-	.001342	.002095	.001906	.000507	.000997		
19	-	.002095	.000000	.001168	.000527			
20	-	.002754	.001154	.000525	.000555			
21	-	.003333	.002137	.001832				
22	-	.003846	.000537	.001206				
23	-	.004302	.001516					
24	-	.004709	.002365					
25	-	.000148						
26	-	.000805						

Table III ( $\alpha = .01$ )

m-n	n			
	9	10	11	12
0	.000621	.000728	.000385	.001364
1	.001394	.000168	.000603	.000608
2	.002052	.001550	.000821	.001333
3	.000807	.001112	.001032	.000705
4	.001506	.000727	.001235	.000126
5	.000434	.000394	.000087	.000844
6	.001130	.000100	.000359	.000344
7	.000196	.001238	.000612	.000998
8	.000869	.000972	.000845	.000557
9	.000024	.000731	.001061	
10	.000682	.000513		
11	.001245			
12	.000546			

$\delta = \gamma$  where  $t$  is the largest integer such that  $\gamma = P(U \leq t | H_0) \leq .025$   
and  $\delta = P(U \leq t + 1 | H_0)$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

m-n	n							
	1	2	3	4	5	6	7	8
0	-	-	-	.010714	.009127	.004437	.006061	.000058
1	-	-	-	.009127	.009848	.007517	.004953	.001800
2	-	-	.007143	.005952	.001010	.003688	.004108	.003290
3	-	-	.001190	.003788	.002467	.000225	.003455	.004559
4	-	-	.008333	.000758	.004021	.004021	.002941	.001328
5	-	-	.000758	.008217	.005020	.001164	.002534	.002724
6	-	.002778	.006818	.007018	.005998	.004261	.002219	.003925
7	-	.006818	.000524	.005220	.000760	.001891	.001952	.001348
8	-	.009848	.005769	.004121	.002007	.004618	.001739	.002589
9	-	.012179	.000824	.002731	.003152	.002573	.001562	.000263
10	-	.003022	.005357	.001797	.004102	.000636	.001415	.001506
11	-	.005952	.001471	.000748	.004950	.003117	.001291	.002604
12	-	.008333	.005392	.000026	.001153	.001404	.001189	.000626
13	-	.010294	.001264	.004783	.002146	.003611	.001101	.001715
14	-	.011928	.004825	.003811	.003049	.002071	.001027	.002689
15	-	.001608	.001692	.003091	.003826	.000592	.000964	
16	-	.003947	.004870	.002320	.000676	.002621	.000910	
17	-	.005952	.001849	.001680	.001576	.001273		
18	-	.007684	.004743	.000987	.002371	.003106		
19	-	.001285	.001957	.000442	.003098			
20	-	.003261	.004615	.003950	.000398			
21	-	.005000	.002094	.003317				
22	-	.006539	.004548	.002778				
23	-	.007906	.002285					
24	-	.001190	.004557					
25	-	.002833						
26	-	.004310						

m-n	n			
	9	10	11	12
0	.005008	.003371	.001346	.002549
1	.003263	.003518	.003070	.002144
2	.001774	.003715	.001276	.001823
3	.000464	.000335	.002886	.001567
4	.003312	.000798	.001369	.001362
5	.002208	.001238	.002864	.001198
6	.001208	.001646	.001542	.001067
7	.000308	.002031	.000304	.000963
8	.002717	.002390	.001749	.000880
9	.001868	.000022	.000641	
10	.001150	.000484		
11	.000455			
12	.002507			



Table III ( $\alpha = .05$ )

$\delta = \gamma$  where  $t$  is the largest integer such that  $\gamma = P(U \leq t | H_0) \leq .05$   
and  $\delta = P(U \leq t + 1 | H_0)$  for  $m + n \leq 30$  and  $\text{Min.}(m, n) \leq 12$ .

m-n	1	2	3	4	5	6	7	8
0	-	-	.000000	.021429	.002381	.003463	.001340	.008508
1	-	-	.021429	.018254	.008874	.013288	.003069	.003640
2	-	-	.014286	.016667	.013384	.009374	.004633	.008430
3	-	.002381	.002381	.013636	.003380	.006044	.006088	.004579
4	-	.014286	.016667	.013636	.008541	.003297	.007328	.001060
5	-	.022222	.007576	.012238	.000383	.000873	.008442	.005477
6	-	.005556	.018182	.012038	.004899	.008468	.001600	.002491
7	-	.013636	.011538	.011172	.009114	.006287	.003113	.006337
8	-	.019697	.006044	.010989	.002731	.004334	.004458	.003699
9	-	.024359	.014835	.010504	.006484	.000870	.005653	.001254
10	-	.006044	.008929	.010458	.000851	.000987	.000255	.004752
11	-	.011905	.004412	.010010	.004494	.005313	.001650	.002543
12	-	.000000	.012010	.000258	.007545	.005236	.002901	.000469
13	-	.005882	.007688	.000710	.002837	.003840	.004034	.003649
14	-	.010784	.003509	.000923	.005816	.002547	.005062	.001744
15	-	.014912	.010150	.001327	.001515	.001346	.000805	
16	-	.002632	.006494	.001534	.004348	.000226	.001944	
17	-	.007143	.003134	.001858	.000390	.004836		
18	.000000	.011039	.008992	.002040	.003114	.003737		
19	.002381	.002569	.005652	.002308	.005522			
20	.004545	.006522	.002692	.002479	.002037			
21	.006522	.010000	.007949	.002718				
22	.008333	.000769	.005128	.002855				
23	.010000	.004416	.002381					
24	.011538	.007672	.007143					
25	.012963	.000739						
26	.014286	.004023						

m-n	n			
	9	10	11	12
0	.003044	.005395	.006027	.005633
1	.002640	.000691	.029159	.001179
2	.002411	.003455	.003346	.002494
3	.002274	.005853	.002287	.003695
4	.002221	.002179	.001369	.000049
5	.002213	.004514	.000567	.001365
6	.002246	.001293	.004689	.002567
7	.002301	.003539	.003901	.003668
8	.002377	.000668	.003193	.000728
9	.002396	.002804	.002552	
10	.002559	.000212		
11	.002659			
12	.002763			

m-n	n
	1
27	.015517
28	.016667

- (1) Aule, ponovan. "Extended tables for the Mann-Whitney statistic," Bulletin of the Institute of Educational Research, Indiana University, Vol. 1, No. 2 (1953), 39 pp.
- (2) Barton, D. E., and David, F. N. "A test for birth order effect," Annals of Human Genetics, 22 (1958), 250-257.
- (3) Barton, D. E., David, F. N., and Mallows. "Non-randomness in a sequence of two alternatives. I. Wilcoxon's and allied test statistics," Biometrika, 45 (1958), 166-177.
- (4) Birnbaum, Z. W. "On a use of the Mann-Whitney statistic," Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Vol. I: Theory of Statistics, editor J. Neyman, Berkeley and Los Angeles: University of California Press, (1956), 13-17.
- (5) Birnbaum, Z. W., and Klase, O. M. "Bounds for the variance of the Mann-Whitney statistic," Annals of Mathematical Statistics, 28 (1957), 933-945.
- (6) van Dantzig, D., "On the consistency and the power of Wilcoxon's two sample test," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 54 (1951), 1-8. (also Indagationes Mathematicae, 13 (1951), 3-10.
- (7) David, F. N. "A note on Wilcoxon's and allied tests," Biometrika, 43 (1956), 485-488.
- (8) Dwass, M. "The large sample power of rank order tests in the two-sample problem," Annals of Mathematical Statistics, 27 (1956), 352-373.
- (9) van Eeden, Constance, and Himilrijke, J. "A test for the equality of probabilities against a class of specified alternative hypothesis, including trend. I and II," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 58 (1955), 191-198, 301-308.
- (10) Festinger, Leon. "The significance of difference between means without reference to the frequency distribution functions," Psychometrika, 11 (1946), 97-105.
- (11) Fix, Eveyln, and Hodges, J. L., Jr. "Significance probabilities of the Wilcoxon test," Annals of Mathematical Statistics, 26 (1955), 301-312.
- (12) Haldane, J. B. S. "The Wilcoxon and related tests of significance," Experientia, 12 (1956), 205.
- (13) ----- and Smith, C. A. B. "A simple exact test for birth-order effect," Annals of Eugenics, 14 (1948), 117-124.

\* The bibliography attempts to be complete.

- (14) Hemelrijk, J, "A family of parameterfree tests for symmetry with respect to a given point," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 53 (1950), 945-955, 1186-1198.
- (15) -----. "Note on Wilcoxon's two sample test when ties are present," Annals of Mathematical Statistics, 23 (1952), 133-135.
- (16) -----. "Exemple d'application des methodes non-parametriques et un nouveau test pour l'egalite de plusieurs probabilities," Report sp 38 of Mathematical Centre, Amsterdam, (1954).
- (17) Hodges, J. L., Jr., and Lehmann, E. L. "The efficiency of some nonparametric competitors of the t-test," Annals of Mathematical Statistics, 27 (1956), 324-335.
- (18) Hoeffding, F. "A class of statistics with asymptotically normal distributions," Annals of Mathematical Statistics, 19 (1948), 293-325.
- (19) -----. "The large-sample power of tests based on permutations of observations," Annals of Mathematical Statistics, 23 (1952), 169-192.
- (20) Kemperman, J. H. De verdelingsfuncties van het aantal inversies in de test van Mann en Whitney, Rapport T. T. no. 7 van de Afdeling Toegepaste Wiskunde, Mathematische Centrum, Amsterdam, (1950).
- (21) Kruskal, W. H. "A nonparametric test for the several-sample problems," Annals of Mathematical Statistics, 23 (1952), 525-540.
- (22) -----. "Historical notes on the Wilcoxon unpaired two-sample test," Journal of the American Statistical Association, 52 (1957), 356-360.
- (23) -----, and Wallis, W. A. "Use of ranks in one-criterion variance analysis," Journal of the American Statistical Association, 47 (1952), 583-621 and 48 (1953), 907-911.
- (24) Lehmann, E. L. "Consistency and unbiasedness of certain non-parametric tests," Annals of Mathematical Statistics, 22 (1951), 165-179.
- (25) -----. "The power of rank tests," Annals of Mathematical Statistics, 24 (1953), 23-43.
- (26) -----, and Stein, C. "On the theory of some non-parametric hypothesis," Annals of Mathematical Statistics, 20 (1949), 28-45.
- (27) Mann, H. B., and Whitney, D. R. "On a test of whether one of two random variables is stochastically larger than the other," Annals of Mathematical Statistics, 18 (1947), 50-60.

- (28) Mood, A. M. "On the asymptotic efficiency of certain non-parametric two-sample tests," Annals of Mathematical Statistics, 25 (1954), 514-522.
- (29) Natrella. "Critical values for Wilcoxon rank sum test," (1954), unpublished.
- (30) Pitman, E. J. G. Lecture notes on nonparametric statistical inference, New York: Columbia University, (1949).
- (31) Probability tables for the Wilcoxon test when there are ties, National Bureau of Standards Report No. 1859, Washington, D. C.
- (32) Putter, J. "The treatment of ties in some nonparametric tests," Annals of Mathematical Statistics, 26 (1955), 368-386.
- (33) van der Reyden, D. "A simple statistical test," Rhodesia Agricultural Journal, 49 (1952), 96-104.
- (34) Rijkoort, P. J. "A Generalization of Wilcoxon's test," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 55 (1952), 394-404 and 56 (1953), 407.
- (35) -----, and Wise, M. E. "Simple approximation and nomograms for two ranking tests," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 56 (1953), 295-302.
- (36) Ruist, E. "Comparison of tests for non-parametric hypothesis," Arkiv For Matematik, Band 3 nr 9 (1954), 133-163.
- (37) Smith, L. J. "On the distribution of the test statistic of Kendall and Wilcoxon's two sample test when ties are present," Neerlandia Statistica, 10 (1956), 205-214.
- (38) Stoker, D. J. "An upper bound for the deviation between the distribution of Wilcoxon's test statistic for the two-sample problem and its limiting normal distribution for finite samples. I," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 57 (1954), 599-606.
- (39) -----, "An upper bound for the deviation between the distribution of Wilcoxon's test statistic for the two-sample problem and its limiting normal distribution for finite sample. II," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 57 (1954), 607-614.
- (40) Sundrum, R. M. "The power of Wilcoxon's two-sample test," Journal of the Royal Statistical Society (B), 15 (1953), 246-252.
- (41) -----, "A further approximation to the distribution of Wilcoxon's statistic in the general case," (1953), unpublished.
- (42) Terpstra, T. J. "A non-parametric test for the problem of k samples," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 57 (1954), 504-512.

- (43) ----- . A non-parametric k-sample test and its connection with the H-test, Report S92 (VP<sub>2</sub>), Mathematisch Centrum, Amsterdam, (1952), 11pp.
- (44) van der Vaart, H. R. "Some remarks on the power function of Wilcoxon's test for the problem of two samples. I," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 53 (1950), 494-506. (also Indagationes Mathematicae, 12 (1950), 146-158.)
- (45) ----- . "Some remarks on the power function of Wilcoxon's test for the problem of two samples. II," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 53 (1950), 507-520. (also Indagationes Mathematicae, 12 (1950), 159-172.)
- (46) ----- . Gebruiksaanwijzing voor de tests van Wilcoxon, Rapport S32 (M4) van de Statistische Afdeling van het Mathematische Centrum, Amsterdam, (1950).
- (47) ----- . "An investigation on the power function of Wilcoxon's two sample test if the underlying distributions are not normal," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 56 (1953), 438-448. (also Indagationes Mathematicae, 15 (1953).
- (48) ----- . "A closed expression for certain probabilities in Wilcoxon's two sample test," Experientia, 12 (1956), 14-15.
- (49) ----- . "Some formulae for the exact computation of certain probabilities in Wilcoxon's two sample test," (1958), unpublished.
- (50) van der Waerden, B. L. "Order tests for the two-sample problem and their power," Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 55 (1952), 453-458 and 56 (1953), 80.
- (51) ----- . "Order tests for the two sample problem," ( second and third communication ), Proceedings Koninklijke Nederlandse Akademie van Wetenschappen (A), 56 (1953), 303-310 and 311-316.
- (52) White, C. "The use of ranks in a test of significance for comparing two treatments," Biometrics, 8 (1952), 33-41.
- (53) Whitney, D. R. A comparison of the power of non-parametric tests and tests based on the normal distribution under non-normal alternatives. Doctor's Thesis, The Ohio State University, 1948, 57pp., typed.
- (54) ----- . "A bivariate extension of the U statistic," Annals of Mathematical Statistics, 22 (1951), 274-282.
- (55) Wilcoxon, F. "Individual comparison by ranking methods," Biometrics, 1 (1945), 80-83.

- (56) ----- "Probability tables for individual comparisons  
by ranking methods," Biometrics, 3 (1947), 119-122.
- (57) ----- "Some rapid approximate statistical procedures,"  
Stanford Research Laboratories, American Cyanamid Company,  
Stanford, Connecticut revised 1949. (also Annals of the New  
York Academy of Sciences, 52 (1950), 808-814.)

Department of Statistics  
University of Minnesota

BASIC DISTRIBUTION LIST FOR UNCLASSIFIED TECHNICAL REPORTS

for contract Nonr 2582(00), Task NR 042-200

<u>ADDRESS</u>		<u>ADDRESS</u>	
Head, Statistics Branch Office of Naval Research Washington 25, D. C.	2	Professor W.G. Cochran Department of Statistics Harvard University Cambridge, Massachusetts	1
Commanding Officer Office of Naval Research Branch Office Navy Nr. 100 Fleet Post Office New York, New York	2	Professor Benjamin Epstein Applied Mathematics and Statistics Lab. Stanford University Stanford, California	1
ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia	10	Professor W. Hirsch Institute of Mathematical Sciences New York University New York 3, New York	1
Office of Technical Services Department of Commerce Washington 25, D. C.	1	Dr. Paul G. Hoel Department of Mathematics University of California Los Angeles 24, California	1
Technical Information Officer Naval Research Laboratory Washington 25, D. C.	6	Professor Harold Hotelling Associate Director Institute of Statistics University of North Carolina Chapel Hill, North Carolina	1
Professor T. W. Anderson Department of Mathematical Statistics Columbia University New York 27, New York	1	Professor L. Hurwicz School of Business Administration University of Minnesota Minneapolis 14, Minnesota	1
Professor Z. W. Birnbaum Laboratory of Statistical Research Department of Mathematics University of Washington Seattle 5, Washington	1	Professor Leo Katz Department of Statistics Michigan State University East Lansing, Michigan	1
Professor A. H. Bowker Applied Mathematics and Statistics Lab. Stanford University Stanford, California	1	Professor Oscar Kempthorne Statistics Laboratory Iowa State College Ames, Iowa	1
Professor Ralph A. Bradley Department of Statistics and Statistical Laboratory Virginia Polytechnic Institute Blacksburg, Virginia	1	Dr. Carl F. Kossack Statistical Laboratory Engineering Administration Building Purdue University Lafayette, Indiana	1
Professor Herman Chernoff Applied Mathematics and Statistics Lab. Stanford University Stanford, California	1		



ADDRESS

Professor Gerald J. Lieberman  
Applied Mathematics and Statistics  
Laboratory  
Stanford University  
Stanford, California 1

Professor William G. Madow  
Department of Statistics  
Stanford University  
Stanford, California 1

Professor J. Neyman  
Department of Statistics  
University of California  
Berkeley 4, California 1

Professor Herbert Robbins  
Mathematical Statistics Department  
Columbia University  
New York 27, New York 1

Professor Murray Rosenblatt  
Department of Mathematics  
Indiana University  
Bloomington, Indiana 1

Professor L. J. Savage  
Statistical Research Laboratory  
Chicago University  
Chicago 37, Illinois 1

Dr. Herbert Solomon  
Teachers College  
Department of Statistics  
Columbia University  
New York 27, New York 1

Professor Frank Spitzer  
Department of Mathematics  
University of Minnesota  
Minneapolis 14, Minnesota 1

Professor S. S. Wilks  
Department of Mathematics  
Princeton University  
Princeton, New Jersey 1

ADDRESS

Professor Evan J. Williams  
Institute of Statistics  
State College Section  
North Carolina State College  
Raleigh, North Carolina 1

Professor J. Wolfowitz  
Department of Mathematics  
Cornell University  
Ithaca, New York 1